





What's Inside

This flipbook has three sections:

TB Testing
Panel A1**Latent TB Infection**
Panel B1**TB Disease**
Panel C1

Each section covers basic information from the other sections, so it is only necessary to use the one section that corresponds to the educational needs of your patient.

How to use this flipbook: Instructions for the health care professional



This flipbook is intended to be a tool to assist the TB patient education process with English-speaking TB patients and their family members. Sitting across from one and other, or side by side, black and white panels such as this one are to be viewed by the health care professional, while color panels are to be viewed by the patient.

Health care professionals are encouraged to read the text on the black and white provider panels to the patient, taking care to tailor the information to the needs of each patient as appropriate. Although this language may seem very basic, it was carefully developed according to health literacy principles and closely follows the language used in the Centers for Disease Control and Prevention TB patient education material series, which is available in English, Spanish, Vietnamese and Tagalog. These corresponding patient materials can be ordered from CDC at: Online:

http://wwwn.cdc.gov/pubs/CDCInfoOnDemand.aspx?Pubid=221235 Mail:

http://www.cdc.gov/tb/publications/pdf/orderform.pdf

It is recommended that appropriate materials from this series be provided to the patient to reinforce the patient education process.





TB Testing



What are the tests for TB?

There are two different tests that are used to see if you have ever had TB germs in your body.

- The TB skin test.
- The TB blood test.

Your healthcare provider will talk to you about which test you will have.

What are the tests for TB?





- There are **two tests** that can tell
 - whether you have ever had TB germs in your body. These tests are used to help diagnose TB infection and TB disease.
- The two tests are the TB skin test and the TB blood test.
- Both the skin test and blood test can be used to test for TB, but the tests work differently. Sometimes one test can't be used, such as if you have you have another medical condition, or if one test is not available in your area.

What is the B skin test?

- A very small needle is used to place a harmless fluid under the skin on the inside of your arm.
- You will feel only a light pinch.

What is the TB skin test?





- A very small needle is used to place a harmless fluid under the skin on the inside of your arm.
- You will feel only a **light** pinch.

How do lease my TB skin test results?

Come back in 2-3 days.

- Any bump that may appear on your arm will be measured.
- Only a health care provider can read your result correctly.



A.3

How do I get my TB skin test results?

- You must come back to the office 2-3 days after the fluid is placed.
- Your health care provider will tell you
 exactly when to come back to the office to have your skin test result read.
- At that time, a health care provider will look at the test spot and measure any bump that may appear there.
- A health care provider is the only person who can read your result correctly. It is important to come back, even if you see nothing on your arm.

What is the B B D D D Est

There are also blood tests that can be used for TB.

- A blood sample will be collected with a needle and tested.
- You only need to come to the clinic once for this test.
- The results are not affected by BCG vaccine.

What is the TB blood test?





Note to health care professional: Skip this panel if your

health care facility is not using this test.

- There are new blood tests used for diagnosing TB Infection.
- Someone in will take a blood sample from you and the blood will be tested.
- With the blood tests (QuantiFERON® -TB Gold In-Tube and T-SPOT® .TB) you will only need to come to the clinic once.
- Some people are concerned that if they have had a BCG vaccine their TB test result will be positive. The BCG vaccine for TB (which is used in many countries, but is not used in the United States) does NOT affect the results of the TB blood test.

What does my **TB test result mean?**

If your test is:	
Positive	Negative
	You don't have TB germs in your body, or
You have TB germs in your body, or	TB germs are not showing up in your body at this time, or
Yo <mark>u had TB germs</mark> in your body in the past.	The test is not able to find the TB germs in your body at this time, or
	Your body is not able to react to the test at this time.



A.5

What does my TB test result mean?

- If your TB test is **positive**, it means you currently have TB germs in your body or you have had TB germs in your body in the past.
- If your TB test is **negative**, it means you do not have TB germs in your body, or the test is not able to find the TB germs in your body at this time, or your body is not able to react to the test at this time.

When-your TB test is **DOSITIVE**.

- You will need more tests to see if you have TB infection or TB disease.
- You will be asked to cough up a sample of phlegm.
- You will also need a chest X-ray.
- You may need other tests.



You will need more tests to see if you have TB infection or TB You will be asked to cough up a sample of phlegm. You will also need a chest X-ray. ou may need other tests



When your TB test is positive...

- If your TB test is positive, you will need more tests to see if you have TB infection or TB disease. This is important because the treatment will be different.
- You will be asked to cough up a sample of phlegm. A health care provider at the clinic will provide you a container and instructions on how to do this.
- You will also need an X-ray of your chest. In some cases, your doctor may also ask for other tests, if he or she feels that you need them.

When your IB test is negative...

You still may need to have more tests if:

- You have been around someone with TB disease.
- Your TB test was done too close to the time you were infected.
- You have signs of TB disease.
- You have a condition that makes it hard for your body to react to the test.







When your TB test is negative...

Sometimes, even when your TB test is negative, you may need to have more tests to find out if you have TB infection or TB disease. You still may need to have more tests if:

- You have been around someone with TB disease.
- Your TB test was within 2 weeks of your exposure to TB.
- You have signs of TB disease like coughing, chest pain, fever, weight loss, or tiredness.
- You have a condition that makes it hard for your body to react to the test.

TB Infection

TB Infection



What is TB?



TB Infection

- Dormant (sleeping) TB germs in your body.
- Will not feel sick.
- CANNOT pass these germs to others.
- Need to take TB medicine to kill TB germs in your body.

TB disease



- Active TB germs in your body.
- May feel sick.
- CAN pass TB germs to others.
- Need to take TB medicine for at least 6 months to be cured.



What is TB?





TB is a short name for tuberculosis. **TB** is a disease that usually affects the lungs, but sometimes affects other parts of the body. I am going to briefly describe TB infection and disease, and then discuss TB infection in more detail.

- **TB infection** means you have only dormant (inactive) TB germs in your body.
- This means the TB germs are sleeping, so they are not making you sick and you cannot pass them to anyone else.
- If you don't take medicine to kill the TB germs now, you can get sick with TB disease in the future.
- If you get sick with TB disease, you can pass TB germs to others.
- If you have TB infection you will need to take TB medicine to kill the TB germs in your body.
- **TB disease** means you have active TB germs in your body that can make you and others around you very sick.
- You may: cough, lose weight, feel tired, have a fever, chills, or have night sweats until you get treatment.
- If you have TB disease you will need to take several TB medicines correctly for at least 6 months to be cured.

What is TB infection?

- You have dormant (sleeping)
 TB germs in your body.
 - Sleeping TB germs will not make you feel sick.
- You CANNOT pass these TB germs to others.
- If the TB germs wake-up and multiply, you will get sick with TB disease.



What is **TB** infection?

- Just to review the last panel, TB infection means you have only dormant (inactive)TB germs in your body.
- One way to understand this is to think of the TB germs as "sleeping".
- Since the TB germs are asleep, they are not making you sick and you cannot pass these TB germs to anyone else.
- If the TB germs "wake-up" and multiply, you will get sick with TB disease.

How did get TB infection?

Spread through the air.

- If you breathe in TB germs, you may get TB infection.
- If the TB germs wake-up and multiply, you will get sick with TB disease.

TB medicine is the only way to kill TB germs.



B.3

How did I get TB infection?

- TB germs are spread through the air when a person with TB disease (point to figure with TB disease on patient panel) coughs, laughs, sings, or sneezes.
- If you breathe in TB germs, you may get **TB infection**. This means you have dormant (sleeping), TB germs in your body.
- Treatment with TB medicine is the only way to kill TB germs.

But why should I take medicine if I don't feel sick?

- TB germs are very strong, even when they are dormant (sleeping).
- Sleeping TB germs can wake-up and make you sick.
- The only way to get rid of TB germs is by taking TB medicines.



But why should I take medicine if I don't feel sick?

- Even though the TB germs in your body are dormant (sleeping), **they are very strong**.
- Many germs are killed shortly after you start taking your medicine, but some stay alive in the body for a long time. It takes longer for them to die.
- As long as you have TB germs in your body, they can wake-up, multiply, and make you sick with TB disease.
- The only way to get rid of TB germs is by taking TB medicines.

You are more likely to develop TB disease if you have TB infection and...



- HIV infection
 - or
- Other health problems or
- You have recently been infected with TB germs or
- You were not treated correctly or did not complete treatment for TB infection in the past.



Some people with TB infection are more likely to develop TB disease

Once you have TB infection, you have a higher chance of getting TB disease if you:

• Have HIV infection

When your body is busy fighting against other infections or diseases it may lose some of its ability to keep the TB germs dormant (sleeping).

• Have other health problems

Other health problems may make it hard for your body to fight TB germs.

• Have been recently infected with TB

This is because you will have the most TB germs in your body during the first few years after you have been infected.

• Were not treated correctly for TB in the past If you did not receive the right number of doses of TB medicine or did not take all of your TB medicine in the past, all of the TB germs many not have been killed.

What is the treatment for TB infection?

There are a few different treatment options for TB infection. Your health care provider will talk to you about which medicine is right for you.

- Isoniazid (i-so-ni-a-zid) also called INH is commonly used to treat TB infection.
- Treatment time with INH is usually 9 months.
- **Rifampin** is sometimes used instead.
 - Treatment time with **Rifampin** is **4 months**.





Note to health care professional:

If your patient has a disease or circumstance that requires an alternate treatment plan, please use this opportunity to discuss the alternate plan or to mention that the health care provider will discuss the plan with the patient.

What is the treatment for TB infection?

- There are a few different options used to treat TB infection the right way. Some of the treatments can't be used in some cases, such as if you have another health condition, or are pregnant.
- The treatment plan you receive will be made to meet your specific needs.
- Isoniazid, also called INH is commonly used to treat TB infection.
- INH is usually taken for 9 months to correctly treat and kill all of the TB germs in your body. It can be taken daily or a few times a week.
- Another medicine, Rifampin is sometimes used to treat TB infection.
- Treatment time with Rifampin is 4 months. Rifampin must be taken daily.

What is the treatment for TB infection?

Another treatment option is a combination of two medicines, **Isoniazid** and **Rifapentine**.

- Medicine is taken once a week for 12 weeks (3 months).
- The medicine may be given by Direct Observed Therapy (DOT) which means a health care worker will be with you while you take the medicine every week.

It is important to take this medicine regularly and not miss a dose.





Note to health care professional:

Skip this panel if your facility does not offer the 12-dose Isoniazid-Rifapentine regimen

What is the treatment for TB infection?

- TB infection can also be treated with a combination of Isoniazid and Rifapentine. This treatment is sometimes called the 12-dose regimen.
- Unlike the other treatments, you only need to take Isoniazid and Rifapentine once a week for a total of 12 weeks, or about 3 months.
- Depending on your weight, you may need to take up to 10 pills every week.
- The medicine is given using Direct Observed Therapy (DOT) which means a health care worker will give you your medicine and be with you while you take it. The health care worker will meet you at a place and time that is convenient for you.
- Since you are only taking 12 doses of the medicine, it is important to take it regularly. If you do miss a dose, you and your health care worker can talk about what to do.
- Because the treatment is shorter, some patients prefer this option. Isoniazid-Rifapentine can't be used for all patients though. Your health care provider will talk to you about which medicine is right for you.

Does the medicine have any side effects?

All medicines can have side effects.

Call us right away if you have:

- Nausea, stomach ache or stomach cramps.
- Less appetite or no appetite for food.
- Vomiting.
- Yellow skin or yellow eyes.
- Dark urine.
- Light colored stools.
- Aches or tingling in your toes or fingers.
- A fever and chills.
- Dizziness or lightheadedness.
- Flu-like symptoms.
- Severe tiredness or weakness.
- Skin rash or itching.
- Bruising.
- Easy bleeding when cut or nosebleeds and bleeding from gums.

Most people can take their TB medicines without any problems.




Does the medicine have any side effects?

- Like all medicines, the medicine you take for TB infection can have side effects.
 - If you have:
 - Nausea or stomach aches or cramps, or
 - Less appetite or no appetite for food, or
 - Vomiting, or
 - Yellow skin or eyes, or
 - Dark urine, or
 - Light colored stools, or
 - Aches or tingling in your fingers or toes, or
 - A fever, or
 - Flu-like symptoms, or
 - Dizziness or light-headedness...
- Remember, people react differently to medicines. If you think you are having any reaction to your treatment, let us (*indicate a specific person and/or phone number*) know right away.
- Most people can take their TB medicine without any problems and will not need to be treated ever again.

Take steps to remember to take your pills



Always take your full dose of medicine even though you are feeling good.

Tips:

- Take your medicine at the same time every day.
- Keep medicine in one place.
- Use a pillbox. Put one week's worth of pills in the box.
- Write yourself a note.
- Use a calendar.
- Ask a family member or friend to help you remember.

Take steps to remember to take your full dose of medicine you take you take your full dose of medicine you take your full dose of medicine you take you



Take steps to remember to take your pills

- **Taking medicine every day** can be a challenge. Remind yourself that you are staying on the medicine to kill all the TB germs.
- Remember to always take your full dose of medicine, even though you are feeling good.
- To help you remember to take the medicine, try some of these tips:
 - Take your medicine at the same time every day.
 - Keep your medicine in one place, where you can't miss it.
 - Use a pillbox. Put one week's worth of pills in the box.
 - Write yourself a note and put it someplace you see often, like the bathroom mirror or refrigerator door.
 - Use a calendar to check off the days you have taken your medicine.
 - Ask a family member or friend to help you remember.

B D i s e a s e

M

TB Disease



What is TB?



- Dormant (sleeping) TB germs in your body.
- Will not feel sick.
- CANNOT pass these germs to others.
- Need to take TB medicine to kill TB germs in your body.

TB Disease



- Active TB germs in your body.
- May feel sick.
- CAN pass TB germs to others.
- Need to take TB medicine for at least 6 months to be cured.



What is TB?





TB is a short name for tuberculosis. TB is a disease that usually affects the lungs, but sometimes affects other parts of the body. I am going to briefly describe TB infection and TB disease, and then discuss TB disease in more detail.

- **TB infection** means you have only dormant (inactive) TB germs in your body.
- This means the TB germs are sleeping, so they are not making you sick and you cannot pass them to anyone else.
- If you don't take medicine to kill the TB germs now, you can get sick with TB disease in the future.
- If you get sick with TB disease, you can pass TB germs to others.
- If you have TB infection you will need to take TB medicine to kill the TB germs in your body.
- **TB disease** means you have active TB germs in your body that can make you and others around you very sick.
- You may: cough, lose weight, feel tired, have a fever, chills, or have night sweats until you get treatment.
- If you have TB disease you will need to take several TB medicines correctly for at least 6 months to be cured.

What is TB disease?



- Large number of TB germs in your body.
- Usually in the lungs, but can show up in other parts of the body.
- TB germs are "awake" and may be making you feel sick.
- TB disease can be cured with medicine!

What is TB disease?



Large number of TB germs in your body. Usually in the lungs, but can show up in other parts of the body TB germs are "awake" and

may be making you feel sick. TB disease can be cured with medicine!



- TB disease means you have a large number of TB germs in your body.
- These germs are usually in the lungs, but sometimes can show up in other parts of the body.
- These germs are no longer "sleeping".
- These germs are "awake" and have multiplied, and may be causing you to feel sick.
- TB disease can be cured with medication.

How did I get TB disease?

COUGH COUGH

> TB germs are spread through the air. If you breathe air with TB germs, you may get TB infection. Untreated TB infection can turn into TB disease.

How did I get TB disease?



- How did I get TB disease?
- TB germs are spread through the air from one person to another when a person who is sick with TB disease coughs, laughs, sings or sneezes.
- If you breathe air that has TB germs, you may get TB infection.
- Without treatment, TB infection can turn into TB disease.

What are some Signs Or Symptoms of TB disease?

COUGH

COUGH

COUGH

- A bad cough that lasts three weeks or longer.
- Chest pain.
- Coughing blood or phlegm.
- Weakness or tiredness.
- Having no appetite.
- Chills and fever.
- Sweating at night or during sleep.

Some people may have all or only a few of these symptoms.



What are some signs or symptoms of TB disease?

Some of the signs or symptoms of TB are:

- A bad cough that lasts three weeks or longer
- Pain in the chest.
- Coughing up phlegm or blood from deep inside the lungs.
- Weakness or feeling very tired.
- Having no appetite.
- Chills and fever.
- Sweating at night or when you are sleeping.
- Some people may have all of these symptoms, some people may have very few.

You CANNOT pass TB germs to others by:



- Sharing food.
- Sharing dishes or eating utensils.
- Sharing clothing.
- Smoking or sharing cigarettes with others.
- Saliva shared from kissing.
- Toilet seats.
- Toothbrushes.

The only way to get TB germs in your body is by breathing them in.



C.5

You CANNOT pass TB germs to others by:

- Sharing food.
- Sharing dishes or eating utensils.
- Sharing clothing.
- Smoking, or sharing cigarettes with others.
- Saliva shared from kissing.
- Toilet seats.
- Tooth brushes.

It is not necessary to keep separate dishes or clothing for TB patients.

The **only way** to get TB germs in your body is by breathing them in.

How is TB disease Urgated?



TB disease is usually treated with several medicines.

The most commonly used medicines are:

- Isoniazid (i-so-ni-a-zid)
- Rifampin (ri-fam-pin)
- Pyrazinamide (peer-a-zin-a-mide)
- Ethambutol (eth-am-byoo-tol)

How is TB disease

A PA

TB disease is usually treated with several medicines.

used medicines are: Isoniazid (i-so-ni-a-zid) Rifampin (ri-fam-pin) Pyrazinamide (peer-a-zin-a-mide) Ethambutol (eth-am-byoo-tol)

C.6

How is TB disease treated?

TB disease is treated with a combination of several medicines.

The most commonly used medicines are:

- Isoniazid (i-so-ni-a-zid)
- Rifampin (ri-fam-pin)
- Pyrazinamide (peer-a-zin-a-mide)
- Ethambutol (eth-am-byoo-tol)

How is TB disease Urgated?



 Several medicines, taken together.

- Take the medicines as you are told by your healthcare provider.
- Treatment takes
 at least 6 months.





Note to health care professional: Use this opportunity to provide the patient with any specific instructions for taking TB medicines.

How is TB disease treated?

- You will start by taking several medicines together.
- It is important to take the medicines exactly as you are instructed. This is the only way to kill TB germs.
- You will need to stay on TB medicines for at least 6 months.

Directly Observed Therapy



• DOT

- Will help you through treatment for TB disease.
- Health care worker brings you your pills.
- Can help you stay with your medicine plan.



Directly Observed Therapy



- Directly Observed Therapy, DOT for short, is a program to help you complete your treatment.
- A health care worker will meet you everyday or several times a week to give you your medicine and make sure you don't miss a dose.
- The health care worker will meet you at a place and time that is convenient for you.
- The health care worker can be a great support to you as you complete treatment.

Do the medicines have any side effects?

 Like all medicines, TB medicines can have side effects. Rifampin can turn some body fluids a reddish-orange color.



C.9

Do the medicines have any side effects?

- Like all medicines, the medicines used to cure TB disease can have side effects, but most people can take TB medicines without any problems.
- **For women:** Some birth control pills may not work as well when you take them with medicines for TB.
- A common, harmless, side effect of Rifampin is that it can turn your urine, tears, and saliva a reddish-orange color.
- While this does not cause any discomfort, it may take some getting used to.

Tell us right away if you have:

- A fever or chills.
- A rash or itching.
- Aching joints.
- Aches or tingling in your fingers or toes.
- An upset stomach, nausea, or stomach cramps.
- Vomiting.
- Changes in your eyesight, such as blurred vision.
- Changes in your hearing such as ringing in your ears.
- Dizziness.
- Bruising.
- Easy bleeding when cut.
- Less appetite or no appetite for food.
- Yellow skin or eyes.
- Dark urine.
- Light colored stools.



Tell your healthcare provider right away if you have:

These are the possible side effects that can occur while taking medicines for TB disease. If you have any of these symptoms while taking TB medicines you should tell us (*indicate a specific person and/or phone number*) right away. Most people can take their TB medicines without any problems.

- A fever or chills.
- A rash or itching.
- Aching joints.
- Aches or tingling in your fingers or toes.
- An upset stomach, nausea, or stomach cramps.
- Vomiting.
- Changes in your eyesight, such as blurred vision.
- Changes in your hearing such as ringing in your ears.
- Dizziness.
- Bruising.
- Easy bleeding when cut.
- Less appetite or no appetite for food.
- Yellow skin or eyes.
- Dark urine.
- Light colored stools.

Take your pills the right-way



If you don't take your pills the right way:

- The medicine may lose the power to cure your TB.
- You can stay sick for a longer time.
- You may have to take more medicines that have more side effects.
- You can pass TB germs on to others.



Take your pills the right way

- If you don't take the pills the right way, the TB germs can become resistant to the medicine.
- This means the medicine will have no effect on the TB germs. If this happens:
 - You can stay sick for a longer time, and
 - You may have to take more medicines that have more side effects.
 - Unfortunately, even the new medicines may not cure the TB, and this means you can pass TB germs on to others.

What if I miss a dose of my TB medicine?



- If you are not on DOT and miss one dose ONE time, take the next dose as scheduled.
- If you miss your pills more than one time, call us BEFORE you take the next dose.
- Talk to us about finding a plan that works for you.



What if I miss a dose of my TB medicine?

- If you are not on DOT, and miss one dose or forget to take your pills ONE time, don't worry, just take the next dose as scheduled.
 - If you forget your medicine more than one time, call us (*indicate a specific person and*/ *or phone number*) BEFORE you take the next dose. We will tell you what to do next.
 - If you have trouble remembering your medicine, talk to us (*indicate a specific person and/or phone number*). We will work with you to find a plan that works.

To help fight TB, take these steps:



Take your medicine the right way.

- During the first few weeks of treatment you may be able to pass TB germs to others.
- To protect the health of those around you:
 - Wear a mask if you are asked to, or cover your mouth and nose with a tissue when you cough or sneeze.
 - If you can, allow outdoor air into your living area.
 - Don't spend time in closed spaces with others.

C.13

To help fight TB disease, take these steps:

- Take your medicine the right way as we have discussed. It is very important to take the medicine the right way; otherwise it may be very difficult or not possible to cure your TB.
- At the start of your treatment, you may still be able to pass TB germs to others. For this reason, you may need to wear a special mask for the first weeks of treatment. This will help prevent TB germs from getting into the air and help protect the health of those around you. If you are not wearing a mask, cover your mouth and nose with a tissue when you cough or sneeze.
- Allow as much outdoor air into your living area as possible. This will keep the air moving and make it harder for the TB germs in the air to stay alive.
- Don't spend time in closed spaces with others until your health care provider says you can no longer pass TB germs to others. Again, this is usually after you have been taking TB medicine for a few weeks.
- The health care workers around you (like me) may also wear a special mask to protect themselves.

Talking to family and friends about 13



Let friends and family know:

- Anyone can get TB!
- You are taking the TB medicines the right way.
- Soon, you will no longer be contagious, and in some months you will be cured.
- A health care worker may speak with them.

TB can be prevented treated, and cured!



C.14

Talking to family and friends about TB

Talking about TB with friends and family is not always easy. You may want to share the information you have learned about TB with them. Let them know:

- Anyone can get TB!
- You are taking the TB medicines the right way. This is important because it ensures that soon you will no longer be contagious (able to pass TB germs to others) and in some months your TB will be cured.
- You may also want to let them know that a health care worker may speak with them about their chances of getting TB. They may need to have a TB test and x-ray.
- Also be sure to tell them that TB can be prevented, treated, and cured!

Medicine Tips

Tips to help you remember your TB medicines if you are not on DOT:



- Take your medicine at the same time every day,
- Keep medicine in one place.
- Use a pillbox.
 - Put a week's worth of pills in the box.
- Write yourself a note.
- Use a calendar.
- Ask a family member or friend to help you remember.

Medicine tips

Tips to help you remember your TB medicines if you are not on DOT: Image: State of the s



Note to health care professional: Skip this panel if your patient is or will be on DOT. Taking medicine everyday can be a challenge.

Remind yourself that you are staying on the medicine to kill all the TB germs. Remember to always take your full dose of medicine **even though you are feeling good**. If you are not on DOT, try some of these tips to help you remember to take your medicine:

- Take your medicine at the same time each day.
- Use a pillbox. Put a weeks worth of pills in the box.
- Keep your medicine in one place where you can't miss it.
- Write yourself a note. Put it on the bathroom mirror or on the refrigerator.
- Ask a family member or friend to help you remember.
- Use a calendar to check off the days you have taken your medicine.



RUTGERS Global Tuberculosis Institute

This flipbook was developed by the Global Tuberculosis Institute http://globaltb.njms.rutgers.edu/ (973) 972-0979

Graphic design by Help Design Group www.helpdesigngroup.com info@helpdesigngroup.com

We would like to thank the patients and staff at the following clinics who participated in the field-testing of this flipbook:

Bergen County Department of Health Services Paramus, NJ Bethlehem Health Bureau Bethlehem, PA **Clark County Combined Health District** Springfield, OH Clearwater Health Department Clearwater, FL DeKalb County Board of Health Decatur, Georgia Lattimore Practice Newark, NJ Lyon County Health Department Emporia, KS Montgomery County Health Department Willow Grove, PA Philadelphia TB Control Program Philadelphia, PA **TB** Satellite Clinic Seattle, WA

The Global Tuberculosis Institute at Rutgers, The State University of New Jersey is designated by the Centers for Disease Control and Prevention as a Regional Training and Medical Consultation Center (RTMCC) in the United States. This publication was supported by Cooperative Agreement Number U52PS004090 from the Centers for Disease Control and Prevention. Its contents are solely the responsibility of the authors and do not necessarily represent the official views of the Centers for Disease Control and Prevention.